

Was GRB 980329 at $z \sim 5$?

Andrew S. Fruchter

Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

ABSTRACT

The optical transient (OT) associated with GRB 980329 was remarkably red. It has previously been concluded that this was the result of dust extinction in the host galaxy (Taylor et al. 1998; Reichart et al. 1998; Palazzi et al. 1998). However, an extinction model can only agree with the data if the I band observations taken about 0.8 days after outburst are discounted (Klose, Meusinger, & Lehmann 1998; Reichart et al. 1998); the flux density ratio between the I and the R of a factor ~ 7 is too great to be explained by extinction, given the relatively blue $K - I$ color. Here it is shown that the *entire* observed optical/infrared spectrum is consistent with that which is expected from an unextincted OT at $z \sim 5$. At this redshift, the light in the observer's R band is strongly suppressed by absorption in the $\text{Ly}\alpha$ forest – an effect which has been clearly seen in galaxies in the Hubble Deep Field (Weymann et al. 1998; Spinrad et al. 1998). In spite of its potentially high redshift, GRB 980329 was an unusually bright burst. If GRB 980329 was indeed at $z \sim 5$, and its gamma-rays were radiated isotropically, the implied energy of the burst would be 5×10^{54} ergs. Should GRB 980329 have a host galaxy, deep imaging could confirm or reject the conclusion that this burst was at $z \sim 5$.

1. Introduction

The last eighteen months have seen a dramatic transformation in our understanding of the nature of gamma-ray bursts (GRBs). The discovery of OTs associated with GRBs (*e.g.* van Paradijs et al. 1997; Bond 1997) has led in turn to direct proof that these objects are at cosmological distances (*e.g.* Metzger et al. 1997; Kulkarni et al. 1998). Analyses of the spectra and temporal behavior of the afterglows from the radio to the x-ray (Frail et al. 1997; Wijers, Rees, & Mészáros 1997; Galama et al. 1998) have confirmed that we are observing expanding relativistic fireballs (Goodman 1986; Mészáros & Rees 1997). Nonetheless, the ultimate astrophysical source of GRBs remains obscure.

The most widely discussed mechanisms for producing GRBs – binary neutron-star or black hole - neutron star mergers, and the collapse of massive stars (Eichler et al. 1989;

Narayan, Paczyński, & Piran 1992; Paczyński 1998) – should be associated with star formation, and indeed a number of well-studied GRB hosts do show signs of intensive star-formation (Fruchter et al. 1998; Djorgovski et al. 1998; Bloom et al. 1998). Additionally, the distribution of host galaxy magnitudes is consistent with a model that links the rate of GRBs at a given epoch of time with the star-formation rate at the epoch (Hogg & Fruchter 1998). But it is the location of the GRBs in their host galaxies which one might expect to distinguish between binary merger and massive stellar collapse, also known as hypernovae, models (Paczynski 1998; Livio et al. 1998). Binaries containing a neutron star should travel far from their birthplace before merger due to the kicks neutron stars receive at birth both from the loss of mass in the binary, and from possibly large impulses from the supernovae (Dewey & Cordes 1987; Lyne & Lorimer 1994). Indeed, neutron star binaries may frequently escape their host galaxy before merger (Bloom, Sigurdsson, & Pols 1998), preventing the creation of an afterglow and perhaps even the GRB itself, due to the absence of a dense external working surface (Meszaros & Rees 1993; Sari & Piran 1997). However, GRBs caused by hypernovae would be expected to occur at the locations of star formation, and thus might be frequently enshrouded in dust (Paczynski 1998).

Although the optical colors of some OTs have shown evidence of moderate extinction (Reichart 1998; Kulkarni et al. 1998; Fruchter et al. 1998), and the inability of observers to find OTs for some GRBs has been interpreted as possible evidence of dust obscuration (Groot et al. 1998; Taylor et al. 1998; Paczyński 1998), there is little direct evidence that GRBs occur in regions of high extinction. GRB 980329, however, appeared to be an excellent candidate for a dust enshrouded GRB. Its OT was not discovered until a radio identification (Taylor et al. 1998) allowed optical observers to re-examine images taken on the first night after outburst. The first image to show an OT was an uncalibrated *I*-band image (Klose, Meusinger, & Lehmann 1998). The derived magnitude $I \sim 20$ at 0.8 days after outburst was surprising given that a much deeper *R*-band image taken at the same time (Palazzi et al. 1998) found the source to have $R = 23.6 \pm 0.2$. Although the *I*-band value has now been calibrated and its brightness lowered to $I = 20.8 \pm 0.3$ (this value will appear in a forthcoming version of Reichart et al. 1998) the $R - I$ color, 2.8, is extremely red. Subsequent imaging in *K*, *J* and *R* has shown that this OT also has very red $K - R$ and $J - R$ colors (see Reichart et al. 1998 for a clear and comprehensive review of the observations of this object; a complete table of the optical and near-infrared observations on this object is also available in Palazzi et al. 1998). While the very red $K - R$ and $K - J$ colors can be explained by strong extinction, it is difficult to reproduce the very steep $R - I$ color by such a model, given that the observed $K - I$ is quite blue. Indeed, in order to do so, one must either discount the *I* band image, or stretch the errors to their limits (Reichart et al. 1998). Here, it is assumed that the data are correct and not misleading. To fit the

data, a radical but plausible alternative is proposed: GRB 980329 occurred at a redshift of ~ 5 .

2. Analysis

Although measurements of the OT of GRB 980329 were taken in the K , J , R and I bands, on no single day are good data available in more than two of these bands (see Table 1 of Reichart et al. for a complete list of the available observations). The standard method of dealing with this situation would be to determine the index of the power-law decay of the OT, and use this to interpolate to a common time. However, as will shortly be shown, the crucial observation was the I band detection of the OT, and the object was observed in I only on the first night. We are therefore forced to extrapolate the later time K and J observations back to the first night. To do this, we use two methods, but both rely on the assumption that the flux density in the different bands falls with the same temporal power law. This assumption should be correct so long as a spectral break in the synchrotron emission of the OT did not pass through the waveband of interest between the times of the different observations used for the analysis. As noted by Reichart et al. (1998) the power-law indices obtained for the R -band data, -1.29 ± 0.19 , and the K -band data, -0.98 ± 0.30 , agree within the errors. Since these two bands represent the short and long wavelengths ends of the spectral region of interest, the assumption of a single power-law index should be good for all the data. Furthermore, the average of these two values -1.14 ± 0.15 agrees well with the power-law indices determined for a number of bursts (Fruchter et al. 1998; Pian et al. 1998; Halpern et al. 1998).

If one then employs the average power-law index of -1.14 to extrapolate the J and K magnitudes back to the time of the I and R images on the first night, 29.9 March 1998, one finds estimated magnitudes of $K = 18.7 \pm 0.4$ and $J = 20.4 \pm 0.4$. As a check on these numbers, one can also subtract the observed $R - K$ color determined at the one time when both bands were observed, 2.3 April 1998, from the observed R magnitude for 29.9 March 1998. One then finds a K magnitude of 18.5 ± 0.4 , which is, within the errors, identical to the K magnitude estimated previously. If one additionally adds the $J - K$ color of 6.3 April 1998 to the K magnitude derived for 29.9 March 1998, one again arrives at a J magnitude of 20.4 ± 0.4 . For the remainder of the paper, then, the values of $K = 18.6 \pm 0.4$ and $J = 20.4 \pm 0.4$ will be assumed for 29.9 March 1998. These derived magnitudes have been corrected for Galactic foreground extinction, converted to μJy and plotted in Figure 1. The I band magnitude used in this figure is the recalibrated value ($I = 20.8$) determined by Reichart et al. The foreground Galactic extinction used, $E(B - v) = 0.074$, is that predicted

by the 100 μ IRAS maps following the method of Schlegel, Finkbeiner and Davis (1997).

In addition to the Galactic extinction adjusted flux densities of the OT, the figure displays a sloping line of the form $\nu^{-0.76}$. If, as the temporal index indicates, our observing frequency is above ν_m (the frequency associated with the minimum specific energy γ_m imparted by the shock to the radiating electrons), then the expected (unextincted) spectral slope is 2/3 the temporal index of -1.14 , or 0.76 (Wijers, Rees, & Mészáros 1997). The K , J and I magnitudes clearly fit the spectral slope well, while the R magnitude is about a factor of 7 ± 2 below the line. While it is hard to understand how dust could produce a decrement of a factor of ~ 7 in $R - I$ while leaving $J - I$ unaffected, there is a well-observed astrophysical phenomenon completely consistent with the observations: the absorption of light by the Ly α forest. At high redshift intergalactic clouds of hydrogen cross any random line of sight sufficiently frequently to significantly reduce the observed far UV continuum of a background source. The strength of this absorption depends on the frequency of the light and the redshift of the background source. At redshifts $\lesssim 2$, the absorption primarily occurs at rest wavelengths shorter than the Lyman limit, $912\text{\AA}$; however, at higher redshifts, the Ly α forest becomes sufficiently dense that rest wavelengths up to Ly α ($1216\text{\AA}$) become significantly depressed (Madau 1995). For a $z \sim 5$ object, this depression, or Ly α break, occurs in the observer’s frame at $\sim 7300\text{\AA}$. Thus $z \sim 5$ objects will be R “dropouts”. Indeed, Madau (1995) predicts a suppression of 80% of the flux of an object at $z \sim 5$ shorter than $1216\text{\AA}$, and observations of spectroscopically confirmed $z \sim 5$ galaxies (Weymann et al. 1998; Spinrad et al. 1998) agree well with this prediction (see Figure 4 in Spinrad et al. 1998), with the observed depression perhaps slightly greater than predicted ($\sim 90\%$). Therefore not only does the wavelength of the break in the OT spectrum agree with a $z \sim 5$, but also the magnitude of the break agrees with that observed. It should perhaps also be noted that the “dropout” technique has worked successfully not only at $z \sim 5$, but also at “lower” redshifts ($2 \lesssim z \lesssim 4$) (Steidel et al. 1996; Madau et al. 1996; Lowenthal et al. 1997), and thus is a well-tested technique for obtaining photometric redshifts.

To quantify the range of redshifts consistent with the data, one can convolve a relatively typical CCD response curve (the STIS CCD was used) with the Harris R filter transmission function (KPNO filter #1466) to simulate the full R response of the observing telescopes. One finds that if the Ly α depression is 90%, then the break spectrum would have to be at $\lambda = 7000\text{\AA}$ (*i.e.* $z = 4.75$) to produce a reduction in observed R of 3.5 – still significantly less than that observed. But if the true Ly α decrement is closer to the 80% predicted, the break would have to occur out of the R band and into the I (and thus have a $z > 5.2$) to have the observed colors.

The reader may be concerned that there could be another cause for this large spectral




Fig. 1.— The spectral energy distribution of the OT associated with GRB 980329 estimated for 29.9 March 1998. The *I* and *R* bands were observed at that date; the *K* and *J* bands have been extrapolated from observations at later times (see text). The diagonal line shows the spectral slope expected from synchrotron emission in an expanding fireball with the observed temporal behavior.

break. Decrements nearly this large have been observed due to other causes in very rare objects (*e.g.* the 2800 Å break in the iron low-ionization broad absorption quasar FIRST J1555633.8+351758, Becker et al. 1997). However, the agreement of the break wavelength and the break magnitude would then have to be entirely coincidental. Rather, given the good agreement between break wavelength and expected decrement, the possibility that GRB 980329 occurred at $z \sim 5$ must be taken seriously.

It is important to note that the fireball model does not predict any spectral break of the magnitude of the one seen here (Sari, Piran, & Narayan 1998). And although the lightcurve of GRB 970228 has shown some unpredicted but comparatively small variability (see, for example, Fruchter et al. 1998), in general the evidence that OTs behave according to the fireball model is quite good (Wijers, Rees, & Mészáros 1997; Galama et al. 1998).

There is yet another piece of circumstantial evidence which suggests GRB 980329 may be at $z \sim 5$. The temporal decay of the OT only fits a power-law if one assumes that nearly all of the observed flux in the R band observations is from the OT and thus is not significantly contaminated by the host. Indeed, even a host as faint as $R = 26.5$ would cause a significant distortion of the power-law decay. However, only one other GRB host is this faint (see Hogg and Fruchter 1998 for a review of host magnitudes) – that of GRB 971214. This host has a spectroscopically measured redshift of $z = 3.4$ (Kulkarni et al. 1998) and has $V = 26.5$ (Odewahn et al. 1998). Yet with $z = 3.4$ one would expect the V magnitude to be partially suppressed by $\text{Ly}\alpha$ absorption; thus the source is probably intrinsically brighter. Furthermore, the apparent M_* (the knee of the luminosity function) for galaxies at $z \sim 5$ is expected to be approximately 26 mag (Hogg & Fruchter 1998). The limit on the host magnitude is already fainter than an M_* galaxy at $z \sim 5$.

3. Discussion

Were GRB 980329 truly at $z \sim 5$, the implications for our understanding of the energetics and beaming of GRBs would be profound. As noted by in 't Zand et al. (1998) the fluence of GRB 980329 in the 50-300 keV of 2.6×10^{-5} ergs $\text{s}^{-1} \text{cm}^{-2}$ would place it in the top 4% of GRBs in the Batse 4B catalog (Paciesas et al. 1997; Meegan et al. 1998). With an assumed cosmology of $H_0 = 70$ km/s and $\Omega = 0.3, \Lambda = 0$ this fluence implies an isotropic burst would have emitted $\sim 5 \times 10^{54}$ ergs in gamma rays alone. This is equivalent to the rest mass of a $2 M_\odot$ object, and would therefore imply strong beaming for any of the GRB mechanisms discussed earlier.

However, the implications of a $z \sim 5$ burst may be nearly as great for cosmology

as for the GRB field. GRB 980329 is only one of 9 bursts with well-identified optical transients (see Hogg and Fruchter 1998). This result would imply that $\sim 20\%$ of bursts are at $z > 3$ and $\sim 10\%$ of bursts are at $z \gtrsim 5$. And, if GRB 980239 is any indication, these high-redshift bursts will fairly frequently be bright enough to be detected in the optical by a 1-m telescope! Indeed, the apparent magnitude of these objects could allow unprecedented studies of the galactic and intergalactic medium at extremely high redshifts through high resolution spectroscopy either from 10-m class telescopes on the ground or from NGST.

Although at the time of this writing the OT should have faded by a factor of ~ 500 from the flux densities shown in Figure 1, the opportunity to measure the redshift of GRB 980329 may not be lost. The host galaxies of all other GRBs with confirmed OTs have been found. If GRB 980329 also has a host galaxy, then the light from that host will suffer the same attenuation in the intergalactic medium as the OT, and the $R - I$ color of the host, like that of the OT, should be larger than 2. While quite possible apparent magnitudes of the host ($I \sim 27$, $R \sim 29$) are daunting, they are within the capabilities of the next imaging instrument planned for HST, the Advanced Camera for Surveys. Therefore, the surprising conclusion of this paper, that GRB 980329 may have been at $z \sim 5$, need not go untested.

4. Acknowledgements

I would like to thank Elena Pian, whose comment that the power-law decay of the OT of GRB 980329 implied a very faint (and perhaps as yet undetected) host led me to think carefully about this object. Sylvio Klose, James Rhoads, Daniel Reichart and Francisco Castander generously shared their data and calibrated values of the I-band magnitude before publication. Steve Thorsett provided valuable comments on the draft. I would also like to thank my colleagues at STScI, and, in particular, Mark Dickinson, Harry Ferguson, Piero Madau and Massimo Stiavelli for many enjoyable and informative conversations on the properties of high redshift galaxies.

REFERENCES

- Becker, R. H., Gregg, M. D., Hook, I. M., McMahon, R. G., White, R. L., & Helfand, D. J. 1997, *Astrophys. J.*, 479, L93
- Bloom, J. S., Djorgovski, S. G., Kulkarni, S. R., & Frail, D. A. 1998, *Astrophys. J.*, submitted, astro-ph/9807315

- Bloom, J. S., Sigurdsson, S., & Pols, O. R. 1998, Mon. Not. R. Astr. Soc., submitted, astro-ph/9805222
- Bond, H. E. 1997, IAUC, 6654
- Dewey, R. J., & Cordes, J. M. 1987, Astrophys. J., 321, 780
- Djorgovski, S. G., Kulkarni, S. R., Bloom, J. S., Goodrich, R., Frail, D. A., Piro, L., & Palazzi, E. 1998, Astrophys. J., submitted, astro-ph/9808188
- Eichler, D., Livio, M., Piran, T., & Schramm, D. N. 1989, Nature, 340, 126
- Frail, D. A., Kulkarni, S. R., Nicastro, S. R., Feroci, M., & Taylor, G. B. 1997, Nature, 389, 261
- Fruchter, A. S., et al. 1998, Astrophys. J., submitted, astro-ph/9807295
- Galama, T. J., Wijers, R. A. M. J., Bremer, M., Groot, P. J., Strom, R. G., Kouveliotou, C., & Van Paradijs, J. 1998, Astrophys. J., 500, L97
- Goodman, J. 1986, Astrophys. J., 308, L47
- Groot, P. J., et al. 1998, Astrophys. J., 493, L27
- Halpern, J. P., Thorstensen, J. R., Helfand, D. J., & Costa, E. 1998, Nature, 393, 41
- Hogg, D. W., & Fruchter, A. S. 1998, Astrophys. J., submitted, astro-ph/9807262
- in 't Zand, J. J. M., et al. 1998, Astrophys. J., 505, 119
- Klose, S., Meusinger, H., & Lehmann, H. 1998, IAU Circ, 6864, 2
- Kulkarni, S. R., et al. 1998, Nature, 393, 35
- Livio, M., et al. 1998, in Proceedings of the Fourth Huntsville Gamma-Ray Burst Symposium, ed. C. A. C. A. Meegan, R. D. Preece, & T. M. Koshut (AIP Conference Proceedings, 428), 483
- Lowenthal, J. D., et al. 1997, Astrophys. J., 481, 673
- Lyne, A., & Lorimer, D. 1994, Nature, 369, 127
- Madau, P. 1995, Astrophys. J., 441, 18
- Madau, P., Ferguson, H. C., Dickinson, M. E., Giavalisco, M., Steidel, C. C., & Fruchter, A. S. 1996, Mon. Not. R. Astr. Soc., 283, 1388

- Meegan, C. A., et al. 1998, in *Bamma-Ray Bursts*, ed. C. A. Meegan, R. D. Preece, & T. M. Koshut (Aip Conference Proceedings 428), 3
- Meszaros, P., & Rees, M. J. 1993, *Astrophys. J.*, 405, 278
- Mészáros, P., & Rees, M. J. 1997, *Astrophys. J.*, 476, 232
- Metzger, R. M., Djorgovski, S. G., Kulkarni, S. R., Steidel, C. C., Adelberger, K. L., Frail, D. A., Costa, E., & Frontera, F. 1997, *Nature*, 387, 878
- Narayan, R., Paczyński, B., & Piran, T. 1992, *Astrophys. J.*, 395, L83
- Odewahn, S. C., et al. 1998, *Astrophys. J.*, submitted, astro-ph/9807212
- Paciesas, W. S., et al. 1997, <http://www.batse.msfc.nasa.gov/data/grb/4bcatalog/>
- Paczynski, B. 1998, *Astrophys. J.*, 494, L45
- Palazzi, E., et al. 1998, *Astr. Astrophys.*, 336, L95
- Pian, E., et al. 1998, in *Proceedings of the Fourth Huntsville Gamma-Ray Burst Symposium*, ed. C. A. C. A. Meegan, R. D. Preece, & T. M. Koshut (AIP Conference Proceedings, 428), 504
- Reichart, D. E. 1998, *Astrophys. J.*, 495, L99
- Reichart, D. E., Lamb, D. Q., Metzger, M. R., Quashnock, J. M., Cole, D. M., Castander, F. J., Cooray, A. R., & Vanden Berk, D. E. 1998, *Astrophys. J.*, submitted, astro-ph/9806082
- Sari, R., & Piran, T. 1997, *Astrophys. J.*, 485, 270
- Sari, R., Piran, T., & Narayan, R. 1998, *Astrophys. J.*, 497, L17
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *Astrophys. J.*, 500, 525
- Spinrad, H., Stern, D., Bunker, A., Dey, A., Lanzetta, K., Yahil, A., Pascarelle, S., & Fernandez-Soto, A. 1998, *Astron. J.*, submitted, astro-ph/9809145
- Steidel, C. C., Giavalisco, M., Pettini, M., Dickinson, M., & Adelberger, K. L. 1996, *Astrophys. J.*, 462, L17
- Taylor, G. B., Frail, D. A., Kulkarni, S. R., Shepherd, D. S., Feroci, M., & Frontera, F. 1998, *Astrophys. J.*, 502, L115

van Paradijs, J., et al. 1997, *Nature*, 386, 686

Weymann, R. J., Stern, D., Bunker, A., Spinrad, H., Chaffee, F. H., Thompson, R. I., & Storrie-Lombardi, L. J. 1998, *Astrophys. J.*, accepted, astro-ph/9807208

Wijers, R. A. M. J., Rees, M. J., & Mészáros, P. 1997, *Mon. Not. R. Astr. Soc.*, 288, L51